PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5:		(11) International Publication Number: WO 93/23544
C12N 15/49, 9/10, 15/62 A61K 39/21	A1	(43) International Publication Date: 25 November 1993 (25.11.93)
(21) International Application Number: PCT/GE (22) International Filing Date: 14 May 1993		& Fairweather, 19 Royal Exchange So., Glasgow Gl
(30) Priority data: 9210337.3 14 May 1992 (14.05.92)	C	(81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
(71) Applicant (for all designated States except US): T VERSITY COURT OF THE UNIVERSITY O GOW [GB/GB]; 2 The Square, University Aver gow G12 8QQ (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): SPIBEY, Norm GB]; 51 Speirs Road, Bearsden, Glasgow G61 2 JARRETT, James, Oswald [GB/GB]; 4 Drum nue, Milngavie, Glasgow G62 8NA (GB).	oF GLA nue, Gla nan [GI	I- S- Published S- With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.
A diagramatic representation		ne variable regions in the FeLV gp70 gene
Variable Basicas 4 2 2		
Variable Regions 1 2 3	4	5
gp 70		p15 eny
(57) Abstract		
to the carboxy terminus of VRI. A co-protein such as GS	TAR	d VR5 or effective fragments thereof. VR5 is preferably attached galactosidase may be attached to the amino terminus of the fused achments to the carboxy terminus may reduce the immunogenici-

48 30

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	C8	United Kingdom	NL.	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BC	Bulgaria	HU	Hungary	PL.	Poland
BJ	Benin	Æ	freland	PT	Portugal
BR	Brazil	iπ	lialy	RO	Romania
CA	Canada	JP	Japan -	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic	SD	Sudan
œ	Congo		of Korea	SE	Sweeten
CH	Switzerland	KR	Republic of Korea	SK	Slovak Republic
a	Côte d'Ivoire	KZ	Karakhstan	SN	Senegal
CM	Cameroon	ü	Liuchtenstein	SU	Soviet Union
cs	Czechoslovakia -	LK	Sri Lanka	TD	Chad
Œ	Coxch Republic	1.11	Luxembourg	TC	Togo
₽£	Germany	MC	Munaco	UA	Ukraine
DK	Dunmark	. MG	Madagascar	us	United States of America
ES	Spain	ML.	Mali	VN	
FI	Finland	MN	Mongolia	VN	Viet Nam

FUSION PROTEIN FROM FELINE LEUKAEMIA VIRUS GP70 AS VACCINE

FIELD OF THE INVENTION

The present invention relates to an immunogenic fused recombinant protein derived from feline leukaemia virus (FeLV), and to a protective vaccine comprising the immunogenic protein.

BACKGROUND OF THE INVENTION

FeLV is a retrovirus of the sub-family oncovirinae.

The virus has a simple genomic organisation with three genes, gag, pol and env, encoding viral proteins.

FeLV occurs in three subgroups, A, B or C, which differ in their env genes. FELV-A is present in every isolate and is efficiently transmitted between cats in nature. Viruses of the other two subgroups arise as variants of FeLV-A in individual cats by recombination of endogenous FeLV env genes (FeLV-B) or mutation within the env gene (FeLV-C). Hence the occurrence of subgroups B or C is dependent upon natural infection of cats with FeLV-A.

FeLV is responsible for several fatal diseases in domestic cats including lymphoid and myeloid leukaemias, fibrosarcomas, anaemia, immunodeficiency, enteritis and reproductive failure. The virus is common in nature: a recent survey in the U.K. indicated that the overall

prevalance f infecti n is appr ximately 15% of sick cats and 6% in healthy cats. These cats have a persistent infection with viraemia and are a source of infection for susceptible animals. FeLV is transmitted either by contact through the transfer of infectious saliva or prenatally across the placenta.

Following natural or experimental exposure of cats to FeLV, there are two possible outcomes. Cats either develop a persistent, life-long infection or recover. Persistently infected cats have a viraemia while recovered cats have no virus in their blood. Some cats which recover may harbour a latent infection in the bone marrow for several weeks or months before it is eliminated. Recovered cats are immune to FeLV as evidenced by their resistance to subsequent challenge with virus.

Two factors influence the outcome of infection. The first is the age at which a cat is exposed to virus. Kittens up to 14 weeks of age are very susceptible and essentially all can be affected by virus challenge.

Older kittens are more resistant so that by 16 weeks of age only about 20% can be infected. The other factor governing susceptibility is dose of virus: high doses lead to persistent viraemia while lower does immunise. The consequences of persistent infection are severe and essentially all viraemic cats die within four years of infection. By contrast recovered cats have a normal life span.

Rec very is due to an immune resp nse t the virus. Only antibody responses have been examined in detail. However, it is clear that resistance to reinfection is mediated by virus neutralising antibodies which are directed at the envelope surface glycoprotein the virus, gp70. Attempts have been made to immunise cats against FeLV infection with vaccines containing gp70, some of which have been successful.

European patent specification EP 0377842 describes the production of replication-defective FeLV viral sequences. US patent 4,794,168 relates to the use of immunogens from FeLV envelope protein P15E in vaccines. European patent specifications 0247904, 0173997 and 0216564, International patent publication W085/02625 and US patent 4,789,702 all describe the use of the FeLV envelope protein gp70 or fragments thereof as antigens.

SUMMARY OF THE INVENTION

The present invention is based on the discovery that a fused recombinant protein comprising two regions of the FeLV molecule which are normally non-adjacent, when combined produced an immunogenic determinant capable of inducing anti FeLV antibodies on inoculation into cats. The immunised cats are resistant to challenge with live FeLV.

Thus, the present invention provides an immunogenic fused recombinant protein which comprises:

- a first protein which is feline leukaemia virus (FeLV) gp70 VR1 protein or effective fragment thereof, having fused thereto;
- a second non-adjacent FeLV protein or effective fragment thereof different from the first protein and capable of stabilising the first protein, such that said fused recombinant protein is immunogenic and protective against FeLV infection.

It has been found that the use of gp70 VR1 protein alone, whilst producing antibodies, is not protective against FeLV infection.

It has however been found that the fusion of a second non-adjacent FeLV protein (or fragment thereof) may result in a fused recombinant protein which is both antigenic and protective against FeLV infection. The second protein is preferably also derived from gp70. In particular, the gp70 VR5 protein (or fragment thereof), which may not by itself produce antibodies, is capable of stabilising the first protein and producing a fused protein which surprisingly has immunogenic and protective properties.

Preferably, the second protein is fused to the carboxy terminus of VR1.

Usually, the fused recombinant protein will be free of glycosylation.

The fused recombinant protein, preferably further comprises an antigenic co-protein, such as Glutathione-S-transferase (GST) or beta-galactosidase.

These relatively large co-proteins solubilise the protein and facilitate production and purification thereof.

Moreover, the co-protein may act as an adjuvant in the sense of providing a generalised stimulation of the immune system. Preferably the co-protein is attached to the amino terminus of the fused protein.

Surprisingly, it has been found that co-proteins (such as GST or \$-galactosidase) may be attached to the amino terminus of the fused protein without adversely affecting immunogenicity thereof, and imparting desirable solubility and stability properties. However, it has been found that the attachment of other proteins to the carboxy terminus (whilst being useful in stabilising the fusion protein against degradation during preparation) may adversely affect its immunogenicity. It is postulated that this may be due to conformational changes brought about by the presence of the co-protein attached at the carboxy terminus. Therefore any co-protein attached at the carboxy terminus should be relatively small such as not to disturb the antigenic conformation of the fusion protein.

The present invention also provides a vaccine formulation which comprises the immunogenic fused recombinant protein together with a suitable carrier. Since the protein may be broken down in the stomach, it is preferably administered parenterally (including subcutaneous, intramuscular, intravenous, intradermal etc.

injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use.

The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art.

The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

The present invention also provides a method of constructing a fused recombinant protein which comprises:

- introducing a DNA sequence coding for the first protein into a plasmid;
- opening the plasmid and introducing a DNA sequence coding for the second protein adjacent the first DNA sequence, and
 - expressing the r combinant protein.

In particular, the DNA sequences may be amplified using polymerase chain reaction (PCR). The amplified sequences may then be inserted one after another into adjacent positions in the plasmid. The plasmid may also include a gene coding for GST or similar co-protein.

The plasmid thus constructed may be expressed in any suitable bacteria, such as E. coli.

Whilst the invention has been described with reference to certain named proteins, such as gp70 VR1 and VR5 proteins, it is to be understood that this covers fragments of the naturally occurring protein and similar proteins (for example, having sequence homologies of 50% or greater) with additions, deletions or substitutions which do not substantially affect the immunogenic properties of the fused recombinant protein.

The second protein is one which does not normally lie adjacent to the first protein in the native FeLV gp70 protein. Non-adjacent means that it is not adjacent in the linearised amino acid sequence.

DESCRIPTION OF PREFERRED EMBODIMENTS

Embodiments of the present invention will now be described by way of example only with reference to the following figures and examples.

Figure 1 shows the plasmid pGEX-2T employed in the construction of the fused recombinant protein:

Figure 2 shows a terminal sequence including restriction sites;

Figure 3 shows the native FeLV gp70 VR1 sequence;

Figure 4 shows the native FeLV gp70 VR5 sequence; and

Figure 5 is a diagrammatic representation of the variable regions in the FeLV gp70 gene.

The following examples illustrate the construction and immunogenic properties of the fused recombinant protein GST-VR1-VR5.

Example 1

Construction of a Plasmid Expressing Recombinant gp70 Fusion Protein

The cloning vector used for these experiments is pGEX-2T (Smith & Johnston, Gene 67 (1988) 31-40). The FeLV gp70 gene fragment is cloned into this plasmid to produce a polypeptide fused to the -COOH terminus of the enzyme Glutathione-S-transferase (GST). Previous experiments have demonstrated that large fragments of gp70 (over 20kd) often result in problems of insolubility and degradation. In an attempt to circumvent these problems it was decided to express a range of smaller gp70 fragments. Furthermore, the regions chosen for expression are some of the "variable" regions of gp70, i.e. those regions showing the greatest variation between sub-groups A, B and C.

The region of the FeLV gp70 gene containing VR1 was

amplified using the foll wing primers. The cloned envelope gene was the template used.

- 1) 5'GAG GGA TCC CTA CAT GTT GAC TTA TG 3'
 BamHI
- 2) 5'TAG GAA TTC ACA ATG TGT TCC CTT TG 3'
 ECORI

The PCR product was digested with the restriction enzymes BamHI and EcoRI the sites for which had been incorporated into the primers as shown above. There are no sites for these two restriction enzymes within the amplified fragment.

The digested PCR product was separated, by electrophoresis on an agarose gel, from any contaminating by products of the PCR reaction. The section of the agarose gel containing the DNA fragment was excised and the DNA removed by electroelution. After ethanol precipitation the DNA fragment was cloned into BamHI/EcoRI cut pGEX-2T.

After transformation of a recipient E.coli strain, clones harboring a recombinant pGEX-2T/VR1 plasmid were isolated.

The following primers were used to amplify a DNA fragment spanning the VR5 region:

- 5'GAC GAA TTC CAG GCT TTG TGC AAT AAG ACA CAA 3' ECORI
- 5' TAG GAA TTC GCA TGC GGT GAG TCC AGT GTT ACA 3' EcoRI SphI

Cloned FeLV gp70 gene was again the template used, the PCR product was digested with EcoRI prior to

purification on agarose. This PCR product was cl ned into the pGEX-2T/VRI plasmid described above. Clones carrying VR5 in the correct orientation with respect to VRI were identified both by DNA sequence analysis and restriction enzyme analysis.

Example 2

Preparation of GST-VR1VR5 Recombinant Protein

The glutathione-S-transferase (GST) of Schistosoma japonicum is a small, very soluble enzyme which is tolerant of insertions at its carboxyl end. The gene (Sj26) encoding for this enzyme has been placed under the control of an inducible promoter and introduced into a plasmid together with an ampicillin resistance marker on it. A multiple restriction site has been introduced at the 3' end of Sj26, in tandem with a region coding for a thrombin cleavage site. The resulting plasmid (pGEX-2T) can produce large quantities of a fusion protein, which is extractable using affinity chromatography. The fusion protein is cleavable and the resulting peptide can be purified by repeating the affinity chromatography.

The following materials and methods are part of routine laboratory practice and can be found in "Molecular Cloning, A Laboratory Manual" second edition, Eds.

Sambrook, Fritsch and Maniatis. and also in "Current Protocols in Molecular Biology." Below is a detailed description f the materials and methods used.

Materials
Enriched L Br th
100ml L Broth
1ml 2M glucose
1ml 1M mgCl₂

2000x IPTG
1g isopropyl D thio- galactopyranoside (IPTG)
8ml water make up to 10ml
Filter sterilise, dispense into aliquots of 0.5ml, freeze
at -20°C

1000x PMSF in isopropanol lmg/ml phenyl methyl sulphonyl fluoride

20% TX100
2ml TX100
8ml water
Filter sterilise. Easier to handle at this concentration.

Glutathione/Tris 250ml 1.0M Tris 4.75ml water 30.73mg glutathione Filter sterilise.

a) Preparation of glutathione beads

A 15ml polypropylene tube is taken and 70 mg of dehydrated glutathione agarose beads are swollen with 10 mls of PBS/1% TX100 using gentle agitation for 30 minutes.

They are then precipitated at 2000 rpm for 5 minutes. The beads are resuspended in 5 mls of PBS to wash them, this procedure is repeated three times. (PBS/1%TX100 or PBS/tween are valid alternatives).

- b) Production of Fusion Protein
- 1. Bacterial cells, transfected with a recombinant pGEX plasmid containing the DNA coding for the protein of interest, are cultured in 100 mls of enriched L Broth vernight in the presence of ampicillin.

- 2. The overnight culture is in culated into 900 mls of enriched L Broth (with ampicillin). The bacterial suspension is grown up with vigorous shaking to an 0.D of approximately 0.6 (550nm) (about 2 hours) then IPTG is added to a final concentration of 50ug/ml. The culture is left for a further 90 minutes. Whilst inducing the culture, start to prepare the swollen, washed glutathione beads for use in stage 5 below.
- 3. The culture is then centrifuged at 6000 rpm for 15 minutes. The supernatant is discarded. The pellets are resuspended in 10 mls of PBS/1% TX100. 10ul of PMSF in isopropanol is added.
- 4. The concentrated bacterial suspension is sonicated in ice for 30 seconds and then centrifuged at 18000 rpm for 15 minutes. The supernatant is retained.
- 5. The beads are then resuspended in the supernatant from step 4 above and 20% TX100 added to a final concentration of 1% TX100. The mixture is gently agitated for 1 hour.
- 6. The mixture is centrifuged at 2000 rpm for 15 minutes and the beads washed 3 times as before. One ml of the solution of 20mM glutathione/50mM Tris is added to the beads and left for 30 minutes. The beads are again centrifuged out (the supernatant being retained) and washed once using another lml of the above glutathione solution. The washing is added to the supernatant.

7. The supernatant contains the fusion protein. An aliquot should be retained to verify this. The supernatant is then dialysed overnight in PBS to remove the excess glutathione. It may then be frozen at -200C.

The preparation exhibited a degree of degradation and/or incomplete synthesis. Nevertheless sufficient material from a 2 litre culture was obtained to conduct these experiments (4 mg. total protein).

Example 3

Vaccination

Since young kittens are most at risk of FeLV infection, the immunisation regime is designed to test the capacity of a vaccine to protect kittens at an early age, and to be compatible with the timing of vaccination against other feline viruses.

The route of challenge is by the oronasal route to simulate natural horizontal transmission of FeLV. A dose of virus is used which will cause persistent viraemia in over 90% of 14 week old kittens.

The schedule of vaccination and monitoring is in Table 1. The response of the kittens to vaccination challenge is given in Table 2.

The VR1-VR5 preparation elicited an immune response. A large proportion of the antibodies raised are directed against the GST portion of the recombinant

molecule as measured by ELISA. However, a competition ELISA using native GST and the recombinant protein also indicates that the immunised cats also responded to the FeLV component of the recombinant protein.

Western blot analysis using FeLV as antigen shows that three out of the four VR1-VR5 treated cats (59,61,62) produced easily detectable anti-gp70 antibodies and cat 60 gave a weak response on Western analysis. All controls were negative.

Virus neutralising antibodies were found in one cat prior to challenge and in a further 2 of the 4, four cats 6 weeks post challenge.

Three of the four vaccinated cats (59,61,62) were also found to be negative for virus isolation 3, 6 and 9 weeks post challenge, whilst all 4 control animals developed viraemia.

The cats responses to immunisation and challenge show that the VR1-VR5 preparation is able to produce an anti-gp70 antibody response and protect the animals from virus challenge.

TABLE 1

SCHEDULE OF VACCINATION AND SAMPLING

Age of Kitten	Week from	
(weeks)	challenge	Event
•		
9	- 5	Blood sample
	·	Vaccination-1
12	-2	Blood sample
		Vaccination-2
14	0	Blood sample
		Challenge
•		
17	3	Blood sample
20	6	Blood sample
23	9	Blood sample

TABLE 2 (VACCINATION RESULTS)

GROUP	CAT	£				TIME	(WEE)	KS) RE	LAT	IVE TO	CHA.	TIME (WEEKS) RELATIVE TO CHALLENGE				1.	
	Ŀ	**	5 >	3 M Z	> «	1 2 X	3	>	2 0	33	A	3 N E W	> 4	Z Z Z	4	9	Z
		·			·												
VR-1+5	66 65 9	1111	1111		1111	0000		1 1 1 1	0004	+3++	1+11	1+11	1 + 1 1	+++	1+11	1 + 1	88 7700
CONT	68 07	1111		0000	1111	0000		1 1 1 1	0000	111	++++	++++	++++	+ 1 + 1	++++	++++	0000
A A VIIII	8 8 7 7 7	ntigen in pla isolated fro neutralising IP70 antibody	plas; from sing a body	asma om plasma g antibody y (ELISA) y (Western	a dy) rn blot	t)											

SUBSTITUTE SHEET.

SEQUENCE LISTING (FIGURE 2)

SEQ ID NO

: 1

SEQUENCE TYPE

: Nucleotide with derived protein sequence

SEQUENCE LENGTH : 51 base pairs

STRANDEDNESS

: single

TOPOLOGY

: linear

MOLECULAR TYPE : genomic DNA

ORIGINAL SOURCE

ORGANISM

: Feline leukaemia virus (FeLv)

USE

DEPOSIT

FEATURES

FeLV protein gp70 terminal sequence with

restriction sites

SEQUENCE LISTING (FIGURE 3)

SEQ ID NO : 2

SEQUENCE TYPE : 208 base pairs

STRANDEDNESS

: single

TOPOLOGY

: linear

MOLECULAR TYPE : genomic DNA

ORIGINAL SOURCE

ORGANISM

: Feline leukaemia virus (FeLV)

USE

: as vaccine in fused protein

DEPOSIT

FEATURES

FeLV protein gp70 VR1

SEQUENCE LISTING (FIGURE 4)

SEQ ID NO : 3

SEQUENCE TYPE : Nucleotide with derived protein sequence

SEQUENCE LENGTH : 42 base pairs

STRANDEDNESS : single

TOPOLOGY : linear

MOLECULAR TYPE : genomic DNA

ORIGINAL SOURCE

ORGANISM : Feline leukaemia virus (FeLV)

USE : as vaccine in fused protein

DEPOSIT

FEATURES FeLV protein gp70 VR5

CLAIMS

- 1. An immunogenic fused recombinant protein which comprises:
 - a first protein which is feline leukaemia virus (FeLV) gp70 VR1 protein or effective fragment thereof, having fused thereto;
 - a second non-adjacent FeLV protein or effective fragment thereof different from the first protein and capable of stabilising the first protein, such that said fused recombinant protein is immunogenic and protective against FeLV infection.
- 2. A fused protein according to claim 1 wherein said second protein is fused to the carboxy terminus of said first VR1 protein.
- 3. A fused protein according to any preceding claim wherein the second protein or fragment thereof is derived from FeLV gp70.
- 4. A fused protein according to claim 3 wherein the second protein is FeLV gp70 VR5 protein or effective fragment thereof.

- 5. A fused protein acc rding to any preceding claim which further comprises an antigenic co-protein attached thereto.
- 6. A fused protein according to claim 5 wherein the co-protein is attached to the amino terminus thereof.
- 7. A fused protein according to claim 5 or 6 wherein the co-protein is glutathione -S- transferase or 8-galactosidase.
- 8. A vaccine formulation which comprises the fusion protein of any preceding claim together with a pharmaceutically acceptable carrier.





Fig 2



CCA AAA TCG GAT CTG GTT CCG CGT GGA TCC CCG GGA ATT CAT CGT GAC TGA
Pro Lys Ser Asp Leu Val Pro Arg Gly Ser Pro Gly Ile His Arg Asp ***

Fig 3


3/5

FeLV VR1

Fig 4

FeLV VR5

CAG GCT TTG TGC AAT AAG ACA CAA CAG GGA CAT ACA GGG GCG Gln Ala Leu Cys Asn Lys Thr Gln Gln Gly His Thr Gly Ala CAC TAT CTA GCC GCC CCC AAC GGC ACC TAT TGG GCC TGT AAC His Tyr Leu Ala Ala Pro Asn Gly Thr Tyr Trp Ala Cys Asn ACT GGA CTC ACC Thr Gly Leu Thr



SUBSTITUTE SHEET

International Application No.

L CLASS	IFICATION OF SUBJ	ECT MATTER (If several classificat	tion symbols apply, indicate ath 6	
		t Classification (EPC) or to both Natio		
	. 5 C12N15/4		C12N15/62;	A61K39/21
IL FIELD:	S SEARCHED			
		Minisum Do	currentation Searched	
Classifica	tion System		Classification Symbols	
Int.Cl	. 5	CO7K ; C12N		
		Documentation Searched on to the Extent that such Documents	ther than Minimum Documentation sats are included in the Fields Searched	
M. DOG	MENTS CONSIDERED	D TO BE RELEVANT		
Category o		coment, 11 with indication, where appr	models of the minutes 12	
		with the state of	opinion, or the relevant passages "	Relevant to Claim No.13
A	CORPORAT 2 Decemb cited in	247 904 (CAMBRIDGE BI TION) Per 1987 The application Whole document	OSCIENCE	1
A	WO,A,8 5 RESEARCH 7 Novemb see the & US,A,4	04 871 (SCRIPPS CLIN FOUNDATION) er 1985 whole document 794 168	IC AND	1
•	WO,A,8 5 20 June cited in	the application 02 625 (CETUS CORPOR 1985 the application whole document	ATION)	1
			-/	
"A" document of the control of the c	interes to be or particular for document but publish g date ment which may throw h is cited to establish th too or other special reas ment referring to an or r means ment published prior to r than the priority date of	ral state of the art which is not ar relevance bed on or after the international doubts on priority claim(s) or se publication date of another ton (as specified) al disclosure, usa, exhibition or the international filling date has	"T" later document published after the ar priority date and not in confliction to understand the principle invention. "X" document of particular relevance; cannot be considered novel or car involve an inventive step. "Y" document of particular relevance; cannot be considered to involve a document is combined with one or ments, such combination being of in the art. "A" document member of the same pa	with the application but or theory underlying the the claimed invention anot be considered to the claimed invention in inventive step when the r more other such docu- wious to a person skilled
V. CERTIF				
Pate of the A	ctml Completion of the 27 AUGUS		Date of Mailing of this Internation 2 9 -09-	in Search Report
oternational :	Searching Authority EUR PEAN	PATENT OFFICE	Signature of Authorized Officer CHAMBONNET F.J	
PCT/ISA/II	10 (second short) (Jennery 1	NS)		

International Application N

integery *	CHRUDE OF DOCKMENT, With Indication, where excessed as also also as a company of the company of		
	Citation of Document, with Indication, where appropriate, of the relevant passages	Kalenan	t to Claim No.
ļ			
	EP,A,O 173 997 (CHIRON CORPORATION)	1	
ľ	12 March 1986	-	
	cited in the application see the whole document		
- 1	see the whole document		
ŀ	EP,A,O 293 249 (AMRAD CORPORATION LIMITED)	7	
	30 November 1988	'	
- 1	see the whole document		
P	PROCEEDINGS OF THE NATIONAL ACADEMY OF	1.	
' ·	SCIENCES OF USA.	1	
1	vol. 89, no. 18. September 1992.	İ	
ŀ	WASHINGTON US	ļ	
ĺ	pages 8457 - 8461		
1	BROJATSCH, J. ET AL. 'Feline Leukemia Virus subgroup C phenotype evolves through		
	distinct alterations near the N terminus		
	of the envelope surface glycoprotein'		
ŀ	see the whole document	1	
	PROCEEDINGS OF THE NATIONAL ACADEMY OF		
	SCIENCES OF USA.	1	-
	vol. 81, no. 12, June 1984, WASHINGTON US	Ì	
	AA AY, HO. TE' ORHE TROL' MYRLINGIAN AZ		
	pages 3675 - 3679		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by monoclonal antibodies: localization of a		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by monoclonal antibodies: localization of a determinant of virus neutralization on the		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by monoclonal antibodies: localization of a determinant of virus neutralization on the feline leukemia virus envelope protein gp70'		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by monoclonal antibodies: localization of a determinant of virus neutralization on the feline leukemia virus envelope protein		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by monoclonal antibodies: localization of a determinant of virus neutralization on the feline leukemia virus envelope protein gp70'		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by monoclonal antibodies: localization of a determinant of virus neutralization on the feline leukemia virus envelope protein gp70'		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by monoclonal antibodies: localization of a determinant of virus neutralization on the feline leukemia virus envelope protein gp70'		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by monoclonal antibodies: localization of a determinant of virus neutralization on the feline leukemia virus envelope protein gp70'		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by monoclonal antibodies: localization of a determinant of virus neutralization on the feline leukemia virus envelope protein gp70'		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by monoclonal antibodies: localization of a determinant of virus neutralization on the feline leukemia virus envelope protein gp70'		·
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by monoclonal antibodies: localization of a determinant of virus neutralization on the feline leukemia virus envelope protein gp70'		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by monoclonal antibodies: localization of a determinant of virus neutralization on the feline leukemia virus envelope protein gp70'		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by monoclonal antibodies: localization of a determinant of virus neutralization on the feline leukemia virus envelope protein gp70'		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by monoclonal antibodies: localization of a determinant of virus neutralization on the feline leukemia virus envelope protein gp70'		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by monoclonal antibodies: localization of a determinant of virus neutralization on the feline leukemia virus envelope protein gp70'		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by monoclonal antibodies: localization of a determinant of virus neutralization on the feline leukemia virus envelope protein gp70'		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by monoclonal antibodies: localization of a determinant of virus neutralization on the feline leukemia virus envelope protein gp70'		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by monoclonal antibodies: localization of a determinant of virus neutralization on the feline leukemia virus envelope protein gp70'		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by monoclonal antibodies: localization of a determinant of virus neutralization on the feline leukemia virus envelope protein gp70'		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by monoclonal antibodies: localization of a determinant of virus neutralization on the feline leukemia virus envelope protein gp70'		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by monoclonal antibodies: localization of a determinant of virus neutralization on the feline leukemia virus envelope protein gp70'		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by monoclonal antibodies: localization of a determinant of virus neutralization on the feline leukemia virus envelope protein gp70'		
	pages 3675 - 3679 NUNBERG, J.H. ET AL. 'Method to map antigenic determinants recognized by monoclonal antibodies: localization of a determinant of virus neutralization on the feline leukemia virus envelope protein gp70'		

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.

GB 9300996 SA 74215

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.

The members are as contained in the European Patent Office EDP file on

The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

27/08/93

Patent document cited in search report	Publication date	Pate	ent family ember(s)	Publication date
EP-A-0247904	02-12-87	AU-B-	607427	07-03-91
		AU-A-	7547987	22-12-87
		JP-T-	1503273	09-11-89
************		WO-A-	8707301	03-12-87
WO-A-8504871	07-11-85	US-A-	4663436	05-05-87
		AU-B-	589563	19-10-89
		AU-A-	4354985	15-11-85
		CA-A-	1265648	06-02-90
		EP-A-	0179139	30-04-86
		JP-T-	61502126	25-09-86
		US-A-	4794168	27-12-88
WO-A-8502625	20-06-85	US-A-	4701416	20-10-87
		AU-B-	577557	29-09-88
		AU-A-	3678184	26-06-85
		DE-A-	3484991	02-10-91
		EP-A,B	0164403	18-12-85
		JP-A-	2238884	21-09-90
		JP-T-	61500662	10-04-86
		US-A-	4789702	06-12-88
EP-A-0173997	12-03-86	US-A-	4876089	24-10-89
		US-A-	5152982	06-10-92
EP-A-0293249	30-11-88	AU-B-	607511	07-03-91
		AU-A-	1793288	21-12-88
		MO-Y-	8809372	01-12-88
		DE-A-	3873989	01-10-92
		JP-T-	1503441	22-11-89

OBM Ports

o For more details about this namex : see Official Journal of the European Patent Office, No. 12/82